



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

SOLUTIONS OF PROBLEMS.

ALGEBRA.

403. Proposed by C. N. SCHMALL, New York City.

A torpedo-boat 40 miles from shore strikes a rock, making a rent in her hull which admits water at the rate of 15 tons in 48 minutes. The ship's pumps can expel 12 tons in an hour. If 60 tons of water is sufficient to sink the boat, find the average rate of steaming so that it may reach the shore just as it is about to sink.

SOLUTION BY CHRISTIAN HORNING, Tiffin, O.

If x represents the rate per hour of steaming, then $40/x$ is the number of hours it takes to reach the shore, and in that time $15 \cdot 60 \cdot 40/48x$ tons of water will have entered the hull, and $12 \cdot 40/x$ tons will have been expelled. Hence $(15 \cdot 60/48) \cdot (40/x) - (12/1) \cdot (40/x) = 60$, and $x = 4\frac{1}{2}$.

Also solved by EMMA GIBSON, F. M. MORGAN, HORACE OLSON, CLIFFORD N. MILLS, WALTER C. ELLS, and J. W. CLAWSON.

404. Proposed by V. M. SPUNAR, Chicago, Illinois.

Show that

$$\begin{aligned} (a+b)(a+b-1) \cdots (a+b-n+1) &= a(a-1)(a-2) \cdots (a-n+1) \\ &+ \binom{n}{1} a(a-1)(a-2) \cdots (a-n+2)b + \binom{n}{2} a(a-1)(a-2) \\ &\cdots (a-n+3)b(b-1) + \cdots + b(b-1)(b-2) \cdots (b-n+1). \end{aligned}$$

SOLUTION BY A. M. HARDING, University of Arkansas.

We have

$$(1+x)^a = 1 + ax + \frac{a(a-1)}{2!} x^2 + \frac{a(a-1)(a-2)}{3!} x^3 + \cdots;$$

also

$$(1+x)^b = 1 + bx + \frac{b(b-1)}{2!} x^2 + \frac{b(b-1)(b-2)}{3!} x^3 + \cdots.$$

Multiply and equate the coefficients of x^n . Then

$$\binom{a+b}{n} = \binom{a}{n} + \binom{a}{n-1} \binom{b}{1} + \binom{a}{n-2} \binom{b}{2} + \cdots + \binom{b}{n}.$$

If we multiply both members of this equation by $n!$ we obtain the desired result

Also solved by ELIJAH SWIFT, who proved the proposition by induction.

405. Proposed by E. J. MOULTON, Northwestern University.

Given the alternating series